

Short-term Climate Extremes: Prediction Skill and Predictability

Emily J. Becker¹, Huug van den Dool¹, and Malaquias Peña²

¹Climate Prediction Center, NCEP/NWS/NOAA, MD

²IMSG at Environmental Modeling Center, NCEP/NWS/NOAA, MD

ABSTRACT

Forecasts for extremes in short term climate (monthly means) are examined to understand our current prediction capability and potential predictability. This study focuses on 2 m surface temperature and precipitation extremes over North and South America, and sea-surface temperature extremes in the Niño3.4 and Atlantic hurricane Main Development regions, using the Climate Forecast System (CFS) global climate model, for the period of 1982-2010. The primary skill measures employed are the anomaly correlation (AC) and root-mean-square error (RMSE). The success rate of forecasts is also assessed using contingency tables.

The AC, a signal-to-noise skill measure, is routinely higher for extremes in short-term climate than those when all forecasts are considered. While the RMSE for extremes also rises, especially when skill is inherently low, it is found that the signal rises faster than the noise. Permutation tests confirm that this is not simply an effect of reduced sample size. Both 2 m temperature and precipitation forecasts have higher anomaly correlations in the area of South America than North America; credible skill in precipitation is very low over South America and absent over North America, even for extremes. Anomaly correlations for SST are very high in the Niño3.4 region, especially for extremes, and moderate to high in the Atlantic hurricane Main Development Region. Prediction skill for forecast extremes is similar to skill for observed extremes. Assessment of the potential predictability under perfect-model assumptions finds that predictability and prediction skill have very similar space-time dependence. While prediction skill is higher in CFS version 2 than in CFS version 1, the potential predictability is not.

This work has been published in the *Journal of Climate* in 2012.

Paper published

Becker, E. J., H. van den Dool, and M. Peña, 2012: Short term climate extremes: prediction skill and predictability. *J. Climate*. doi: <http://dx.doi.org/10.1175/JCLI-D-12-00177.1>.




Fig. 1 CFSv2 2 m temperature anomaly correlations over North America, expressed as a function of the target month (horizontal) and lead (vertical), for a) all cases, b) when the model predicts an extreme, defined as ± 1.645 local standard deviation, c) all cases with CV3RE applied (Cross-validation when three years are excluded. See the paper for details.), and d) predicted extremes with CV3RE applied. (From Becker *et al.* 2012)