




NOAA's MAPP-CTB Projects Update: Community R20 Contributions to the Improvement of Operational S2S Climate Prediction

Jiayu Zhou, Climate Mission, Office of Science and Technology Integration
David DeWitt, Climate Prediction Center, National Centers for Environmental Prediction

National Weather Service, National Oceanic and Atmospheric Administration, U.S. Department of Commerce

NOAA's Modeling, Analysis, Predictions and Projections - Climate Test Bed (MAPP-CTB) projects support research to significantly increase the accuracy, reliability, and scope of NWS Climate Prediction Center operational seasonal-to-subseasonal (S2S) climate probabilistic forecast products. Out of twenty-six funded projects, twelve are on track to be completed by the end of fiscal year 2018, seven are in progress with an adjusted deliverable schedule, and seven are new development projects. This presentation will provide stakeholders an update on 1) completed projects leading to an improved operational prediction capability, 2) progress on prominent ongoing projects, and 3) challenging developments supported by leveraging the research community. The CTB management solicits feedback on optimal project performance to accelerate the transition from research to operations, toward the NWS strategic vision of a "Weather-Ready-Nation".




Completed Projects (RL #: 8-9)


System Implementation	Developer
1. Improved Turbulence and Cloud Processes	USU
2. NMME Phase 2	UM-RSMAS
3. NCEP GEFS for Monthly Forecast	NCEP/EMC
4. Real-Time Multi-Model Sub-Seasonal Predictive Capability (SubX)	UM-RSMAS
5. U.S. Monitoring and Prediction System for Flash Droughts	UCLA
6. Ensemble-based Sea Ice Analysis and Forecasting	UMD

System New Component	Developer
7. Lake-Effect Process	USU
8. CCSM4 (SubX)	UM-RSMAS
9. Navy Earth System Model (NESM)	Naval Research

System Analysis and Optimization	Developer
10. NMME Skill, Predictability and Optimum Combination	GMU
11. Assessment of CFS Severe Weather Predictions	CU
12. Estimation of NASA GEOS-5 MJO Forecast Skill and Land Surface Feedback	NASA-USRA



3 Pattern Anomaly Correlation (PAC) for Northern Hemisphere 500 hPa geopotential height for lead weeks 3&4 during the period from May 2014 to May 2016. CTL is in black, SPs in red, SPs+CFSBC in green, SPs+CFSBC+CNV in blue and CFSv2 in brown with period average PAC scores for each configuration (numbers in the bottom of plot with corresponding color). MJO forecast skill (from 12.5 days to 22 days) and NH 500hPa height ensemble mean AC scores (from 0.35 (control) to 0.404) (SubX configuration) are improved after applying the new stochastic physics perturbation, updated SST and new convective schemes. (PI: Y. Zhu)



12 Anomaly correlation coefficient for 5-day averaged H500 anomalies from GEOS-S2S hindcasts when (left column) forecast initialization contains MJO in phases 3 or 7, (middle column) for all forecasts irrespective of MJO signal in initial conditions, and (right column) difference between the two. The rows represent 5-day averaged fields as the forecast progresses from day-16 to day-30. Verification is performed against MERRA-2 data, and forecasts cover the period 1999-2015. All forecasts are initialized over an extended winter period (November through March). The results are promising in that increased skill (the red shading in the right column) is observed up to days 26-30, especially over the eastern US, suggesting a typical MJO-NAO teleconnection pattern. It indicates higher skill of H500 anomalies prediction when an MJO in phases 3 or 7 is present in the initial conditions. (PI: D. Achuthavarier)

Accomplishment Highlights

1. The Simplified Higher-Order Closure (SHOC) and Chikira-Sugiyama-Arakawa-Wu (CSAW) unified cumulus parameterization have been implemented and tested in NEMS/GSM and GFS with FV3 dynamical core (FV3gfs). SHOC, CSAW along with Morrison-Gettelman double moment microphysics have been implemented in the Interoperable Physics Driver 4 of the NEMS/FV3 model. (PI: S. K. Krueger)

2. NMME Partner Agreement was signed by all parties and the Community Earth System Model (CESM) retrospective forecast completed. NMME operational forecasts continue to be delivered on time. A new procedure for estimating forecast spread is developed. (PI: B. Kirtman)

3. A real-time monthly GEFS version with the identical configuration as reforecast but larger ensemble size (21 members) is being conducted every Wednesday since July 2017. Finished a 17-yr reforecast with a high-resolution (34km for 0-8 days and 52km for 8-35 day) model and 11 member ensemble. Eleven real-time priority variables are delivered to CPC. Other priority 2 and 3 variables are delivered to IRI. (PI: Y. Zhu)

4. The SubX began making real-time predictions in July 2017. The data are provided to CPC as guidance to their week 3-4 forecast products and also to IRI for posting on the IRI Data Library. Completed re-forecast database. Comprehensive skill evaluation showed the benefit of the MME over any individual model. (PI: B. Kirtman)

5. An experimental real time flash drought monitor is on the CPC website. Experimental real time flash drought forecast in three categories based on the CFSv2 seasonal forecasts has been implemented. (PI: D. P. Lettenmaier)

6. Sea ice modeling and data assimilation (EnKF) capability has been implemented in CFS, and a full year sea ice analysis using the new ensemble-based system completed. (PI: J. Carton)

7. The 16-year retrospective forecasts with nine leads were performed with CFS and CFS-Flake. These forecasts for the Great Lakes region were quantitatively analyzed with different metrics, showing the predicted surface skin temperature, precipitation, and lake ice spatial distribution with the coupled CFS-Flake were significantly improved. (PI: J. Jin)

8. Real-time Community Climate System Model 4.0 (CCSM4) subseasonal forecasts began in July 2017. All priority 1 variables from all the hindcasts have been provided to IRI. (PI: B. Kirtman)

9. Demonstrated that NESM forecasts can be supplied to NOAA in a timely manner for S2S operational products. Completed the re-forecasting effort from 1999 to start of the real-time experiment, and supplied these outputs to NOAA and IRI. (PI: N. Barton)


10. Developed a new method for determining the ensemble size and initialization frequency of the lagged ensemble that minimized the MSE. Identified the optimal lagged ensemble for monthly, subseasonal and seasonal forecasts in CFSv2. Developed new, rigorous methods (code available online) for comparing forecast skill. (PI: T. Delsole)

11. The automated 00 UTC CFS ensemble mean severe weather guidance dashboard has been transitioned to application (http://www.spc.noaa.gov/exper/CFS_Dashboard/). (PI: M. Tippett)

12. The near real-time production of sub-seasonal forecasts commenced at GMAO, NASA/GSFC on July 25, 2017, and since then, the forecasts (consisting of 10 Priority-1 variables) have been submitted to CPC every Wednesday. In addition, 39 output fields from the forecasts are being submitted to the IRI's SubX data repository. (PI: D. Achuthavarier)




One-Year Extension Projects (RL #: 7-8)

On Finalization	Developer
13. Improved Cloud and Boundary Layer Processes	UW
14. Modeling and Data Infrastructure	UC/CIRES
15. Subseasonal Excessive Heat Outlook System	UMD-ESSIC/CICS
16. S2S Climate Products for Hydrology and Water Management	NCAR
17. Moisture and Snow Data Assimilation in NLDAS	NASA/GSFC
18. Seasonal Forecast Application for AK Wildland Fire Management	UAF
19. Eddy-Permitting Hybrid Global Ocean Data Assimilation System	UMD



15 (a) The dry Excess Heat Factor (EHF) as calculated using ERA-Interim data from 1979-2016 at the closest grid point to Niamey airport, Niger. The reported excessive heat wave, which occurred in April 2010 is well captured. (b) Both dry and wet EHF are successful in capturing the April 2010 event, however, during May as the monsoon season approaches the wet EHF shows higher sensitivity. The wind induced skin evaporation and lower solar incoming radiation provide relief from excessive heat. Further considering margins provided by acclimatization, the geographical modulation of the danger level is going to be explored. The global heat-impact oriented subseasonal excessive heat system runs experimentally on real time at the University of Maryland since May 2018 and provides probabilistic forecasts of the wet and dry EHF. (PI: A. Vintzileos)


Progress on Prominent Ongoing Projects






Challenging Developments Supported by Leveraging the Research Community

New Projects (RL #: 6-7)

Challenges and Prospects	Developer
20. A New Technique for Improved MJO Prediction	NOAA/PMEL
21. Probabilistic Multimodel, Calibrated Subseasonal Global Forecast Products	CU/IRI
22. Sensitivity Analysis of NMME Seasonal Predictions to Ocean Eddy Resolving Coupled Models	UM-RSMAS
23. S2S Prediction Improvement with NCAR's CESM2-WACCM	NCAR/CGD
24. A Hybrid Statistical-Dynamical System for the Seamless Prediction of Daily Extremes and S2S Climate Variability	NOAA/CPC
25. Novel Statistical-Dynamical Forecasts for Tropical S2S Drivers	NCSU/NCICS
26. Predicting Atmospheric Rivers (AR) and Their Impacts in Weeks 2-5 Based on the State of the MJO and QBO	CSU



The figure on left shows the time mean precipitation simulations in mm day⁻¹ with low (contours) and high (shaded) model resolutions (top panel), and the climatological precipitation of CPC Merged Analysis of Precipitation (bottom panel). It demonstrates the presence of resolved ocean eddies modified the mean climate. (PI: B. Kirtman)



Completed Projects Leading to an Improved Operational Prediction Capability